

Supplementary Material

Microbial diversity in a submarine hydrothermal chimney from the serpentinized system of the Prony Bay (New Caledonia) over a 6 years period.

Anne Postec^{1*}, Marianne Quéméneur¹, Méline Bes¹, Nan Mei¹, Fatma Benaïssa¹, Claude Payri², Bernard Pelletier², Christophe Monnin³, Linda Dombrowsky^{1,2}, Bernard Ollivier¹, Emmanuelle Gérard⁵, Céline Pisapia⁵, Martine Gérard⁴, Bénédicte Ménez⁵, Gaël Erauso^{1*}.

1 Aix Marseille Université, CNRS/INSU, IRD, Mediterranean Institute of Oceanography, UM110, 13288 Marseille, France


2 Institut pour la Recherche et le Développement, Centre de Nouméa, promenade Laroque, 98848 Nouméa, Nouvelle-Calédonie

3 Géosciences Environnement Toulouse, UMR 5563, 14 avenue Édouard Belin, 31400 Toulouse

4 Institut de Minéralogie et de Physique des Milieux Condensés, 4 place Jussieu, 75005 Paris, France

5 Institut de Physique du Globe de Paris, Sorbonne Paris Cité, Univ. Paris Diderot, CNRS, 75005 Paris, France

* Correspondence: anne.postec@univ-amu.fr and gael.erauso@univ-amu.fr



Supplementary Figure 6. Phylogenetic tree representing *dsrB* translated gene sequences. The tree was constructed using the neighbor joining method. Bootstrap values <50% are not shown. Clone libraries are distinguished by various markers and colors: 2005 (■ in red), 2010 (▲ in green) and 2011 (● in blue). Pink font denotes phylotypes from other terrestrial alkaline sites, light blue denotes phylotypes from hydrothermal. After each OTU name is indicated the number of clones retrieved for this OTU. Scale bar: 0.05.